



REDLANDS COMMUNITY HOSPITAL

Your Step-by-Step Guide to Hip Replacement


PRE- and POST-Operative Instructions
Exercises & Nutrition

The Hospital Experience
Frequently Asked Questions

Steps to Success

FOR

HIP REPLACEMENT



redlandshospital.org



Your Care TEAM PARTNER

The Navigator

You will have a partner to help you prior to your surgery, during hospitalization and after you return home. This care team partner is called the Navigator. You can reach the Navigator by calling (909) 793-4383.

Your Navigator's name is:

Table of CONTENTS

YOUR CARE TEAM PARTNER	PRE-OPERATIVE CHECKLIST
The Navigator 2	Home checklist 14
Welcome 5	What to bring 15
Steps to Success for Hip Surgery 6	Patient Bill of Rights. 15
	The Navigator 16
PREPARE FOR SURGERY	Health questionnaire 17
Pre-registering with the hospital 7	FOUR WEEKS BEFORE SURGERY
Obtaining medical and anesthesia clearance 7	Start supplements 20
Making an appointment for laboratory tests 7	TEN DAYS BEFORE SURGERY
Required pre-op orientation class 7	Stop medications that increase bleeding 20
Blood transfusions 7	THREE TO FIVE DAYS BEFORE SURGERY
Review "Exercise Your Right". 8	Prepare your home for your return 21
Start pre-op exercises 8	Prescriptions 21
Medications and surgery 8	THE NIGHT BEFORE YOUR SURGERY
Blood pressure medications 8	Do not eat or drink after midnight 22
Diabetes medications 9	What to pack for the hospital 22
Blood thinners 10	Best to leave at home 22
Asthma/Breathing Medications 10	DAY OF SURGERY
Anti-inflammatory Medications 10	Pre-op preparations 23
Vitamins 10	Move to recovery 23
Herbal supplements 10	Surgical waiting area 23
PRE-OPERATIVE CLASS	In your room 24
Introduction 11	One day after surgery 24
Class preview 12	DISCHARGE PLANNING
IMPORTANCE OF YOUR COACH	Discharge day 25
Before surgery 13	HOME CARE
At the hospital 13	General information 26
	Be comfortable 26

Table of CONTENTS

EATING RIGHT FOR FAST RECOVERY

Introduction	27
Healthy eating plan	27
Balanced diet.....	28
Sample menu	29

PAIN CONTROL

Anesthesia during surgery	30
Post-op pain control	30
Prevent the pain cycle	31
Pain scale	31
Other methods to decrease pain	31

PRE-OPERATIVE EXERCISE

Stretch your joint	32
--------------------------	----

HOME CARE

General information	36
Wean yourself from pain medications. ...	36
Prevent constipation	37
Blood clots and anticoagulation.....	37
Care for your incision	38
Medications.....	40
Signs of infection	42

EVERYDAY LIVING

Self-care	45
Bathing	45
Toilet	45
Rest/Sleeping	46
Dressing.....	46
Sitting	46
Standing	46
Kitchen activity.....	47
Laundry	47
Lifting	47
Driving/Riding in a car	47
Preventing weight gain	48

ADVANCED DIRECTIVES

Three types of advanced directives	49
--	----

THE HUMAN HIP

Anatomy of the Hip	50
--------------------------	----

FAQs	51
------------	----

Anesthesia FAQs	58
-----------------------	----

POST-OPERATIVE JOURNAL

My surgery journal	59
--------------------------	----



Welcome to the SPINE & JOINT INSTITUTE at Redlands Community Hospital

You have taken the very first step toward a healthier lifestyle with your decision to have elective hip surgery. Each year, thousands of Americans make the decision to end their chronic hip pain by undergoing hip surgery and are enjoying life again.

We offer a unique program. Our program is designed to return you to an active lifestyle as quickly as possible. Each step of the program is carefully choreographed to give you the best possible results. Here are just a few of the special features you may enjoy:

- Nursing and professional staff dedicated specifically for patients having joint surgery
- Coach's program for your family, spouse or friend
- Casual clothes (no drafty hospital gowns)
- Individual exercise
- Coordinated care after discharge
- Written instructions

*Our team appreciates the opportunity to help you on the road to recovery. You are more than a patient to us; you are **our** patient. We will do our best to make your stay a pleasant one.*

PLEASE NOTE: While you are reading this book, please remember that people are individuals and may vary in terms of their needs.



Important Reminders:

Please call your surgeon's office if you need to reschedule any of the following appointments.

Surgery:

Date: _____

Time: _____

Room: _____

Pre-op Labs:

Tests:	Labs	EKG	Xray
Date:	_____	_____	_____
Time :	_____	_____	_____
Room:	_____	_____	_____

Pre-op Orientation:

Date: _____

Time: _____

Room: _____

My Coach:

Date: _____

Time: _____

Room: _____

Other Instructions to remember:

Steps to Success for HIP SURGERY

Steps to Success for Hip Surgery is designed to be a handy, one-source reference guide that will take you through your upcoming hip surgery and recovery period with ease.

Even an elective surgery such as hip surgery, can be an unsettling prospect. Most patients and their families do not know what to expect. They trust their doctors and nurses, but are still apprehensive about the unknown.

The Spine and Joint Institute team understands your concerns. We will take every step possible to make your surgical experience pleasant and enjoyable. Keeping you informed is key. You will know what to expect and when, and what is considered normal healing. You will also learn what you need to do to hasten your own recovery process.

As you read Steps to Success for Hip Surgery, you will find answers to frequently asked questions, checklists to prepare you for surgery, what to expect during the hospital stay, exercises and other helpful information.



Please follow your healthcare professional's advice in terms of adding to or changing any of the guidelines presented in this book.

prepare for SURGERY

The time before your surgery can be a busy one and it does require a little planning. Use the following list as your guide.

PRE-REGISTERING WITH THE HOSPITAL

Once your surgery is scheduled, you will be contacted by the Navigator to arrange a date and time to confirm your pre-op orientation where you will pre-register for your surgery at the hospital.

OBTAINING YOUR MEDICAL AND ANESTHESIA CLEARANCE FOR SURGERY

Your surgical team will need to be aware of any existing health conditions in order to avoid potential complications during your surgery. If a potential problem is detected, then additional steps will be taken to ensure a successful surgery.

You will need to undergo a brief physical to obtain medical and anesthesia clearance for surgery. Your surgeon's office will need to start this process immediately.

MAKING AN APPOINTMENT FOR LABORATORY TESTS

Your surgeon's office will order any tests needed to be completed prior to surgery.

REQUIRED PRE-OP ORIENTATION CLASS

The pre-op class is a very important part of your preparation for surgery. The class is designed to give you the opportunity to meet our staff, other patients and their families. During this time, you will be given valuable instructions to prepare you for the days to come. Our staff will be available to answer any questions and concerns that you or your family may have. The Navigator from Redlands Community Hospital will contact you with information on the upcoming pre-op orientation class and any lab and EKG tests needed (if ordered).

BLOOD TRANSFUSIONS

During surgery, blood can be lost. It is extremely rare that a blood transfusion is necessary. When a transfusion is needed, the blood is usually supplied by a Blood Bank. You may have blood drawn for a type-and-screen the morning of surgery.

prepare for SURGERY

The time before your surgery can be a busy one and it does require a little planning. Use the following list as your guide.

REVIEW "EXERCISE YOUR RIGHT"

The law requires that everyone being admitted to a medical facility has the opportunity to make advanced directives concerning future decisions regarding their medical care.

Please refer to page 49 for further information. Although you are not required to do so, you may make the directives you desire.

If you have advanced directives, please bring copies to the hospital on the day of surgery.

START PRE-OPERATIVE EXERCISE

Many patients with arthritis favor their joints and consequently become weaker. This can interfere with their recovery. It is important that you begin an exercise program before surgery. Reference page 32 for some joint exercise ideas.

MEDICATIONS AND SURGERY

Your pre-op nurse will confirm with you what and how to take your medications.

Discuss your current medications with your physician to see if and when you should modify your medication schedule. Remember to include any over the counter drugs as well as vitamins and herbal supplements. It is important that your healthcare team know all of the medications, nutritional and herbal supplements that you are currently taking, as they may cause problems such as excessive bleeding during your surgery. This may or may not be relevant to you depending on a number of factors including:

- The surgical procedure you are having
- Your medical history
- The medications/supplements you are taking

Remember, it is very important to provide accurate information to your healthcare team. Do not stop or change any of your medications unless instructed.

BLOOD PRESSURE MEDICATIONS

In general, patients are told to continue taking these medications at the normal scheduled time. Some blood pressure medications are not taken the morning of surgery. Your pre-op nurse will confirm with you what and how to take your blood pressure medications before surgery.

prepare for SURGERY

DIABETES MEDICATIONS

The diabetic medications that you take have a direct effect on your glucose levels, the way your body produces insulin and responds to storing glucose. Each medication is cleared by various organs in the body. Certain medications will be held up to 2 days before surgery, stopped the morning of surgery, or the dosage reduced.

- If you take an oral medication such as Metformin®, you may be told not to take it the day of surgery. Other oral medications such as Glucophage® may be stopped for several days or more prior to surgery as this medication may cause serious side effects while under anesthesia.
- Insulin will be continued, but usually at a different dose in order to prevent your blood sugar from dropping too low during fasting.
- It is important to discuss what your medication regimen should be regarding your diabetes medications prior to surgery with the Navigator and/or your doctor.



This list is by no means exhaustive, so please check with your doctor before taking any of your own medications

*Feel free to ask any questions you may have.
The more you know, the better prepared you will feel.*

prepare for SURGERY

BLOOD THINNERS

Medications such as warfarin and aspirin should be stopped seven to ten days prior to surgery. With some patients, however, the benefits of this medication may outweigh the risks of discontinuing this medication until the day of surgery. This is why your medical history is so important, not just what you are taking, but why. **Please discuss this with your surgeon at least two weeks prior to your scheduled surgery.**

ASTHMA/BREATHING MEDICATIONS

Patients are normally instructed to continue taking these medications according to their regular schedule.

ANTI-INFLAMMATORY MEDICATIONS

These medications usually need to be stopped seven to ten days prior to your scheduled surgery, as they increase the risk of bleeding. These medications include:

- Ibuprofen (Advil, Motrin, Nuprin)
- Naproxen (Anaprox, Naprosyn, Aleve)
- Ketoprofen (Orudis, Oruvail)
- Diclofenac (Voltaren, Cataflam)
- Indomethacin (Indocin)
- Piroxicam (Feldene)

Any other medications that contain aspirin will need to be stopped as well.



This list is by no means exhaustive, so please check with your doctor before taking any of your own medications.



Pre-operative class



The Spine and Joint Institute is not a typical hospital unit. There are no sick patients there, only people who are committed to relieving their spine or joint pain. Treated in a group setting, patients are ready to work towards regaining the quality of life they had before the onset of pain. Therefore, hip patients have their own area and their own set of guidelines which are very different from the rest of the hospital.

Prior to your surgery, you will be attending a pre-op orientation class. This is your time to learn about what to expect over the next few weeks and, more importantly, what will help speed your recovery.

During the required pre-op class, you and your coach will have the opportunity to meet the staff, as well as other patients and their families. Questions will be addressed and other vital information will be discussed.

We highly recommend that you designate your spouse, friend, or family member who will be caring for you after surgery as your official coach. The coach will be at your side, helping you with your exercises, keeping you motivated and generally doing what is necessary to get you back on your feet again and enjoying life. Be sure to invite your coach to attend the pre-op class with you.

Depending upon your progress, you may be going directly home shortly after surgery. Once home, you may need special equipment or training to help with your recovery. The pre-op class is the perfect time for you and your coach to start preparing for your homecoming.

This is your opportunity to learn what is in store for you in the upcoming weeks and what you can do to make your surgery a success. Feel free to ask any questions you may have. The more you know, the better prepared you will feel.

CLASS PREVIEW

- Meet and greet
- Preparing for surgery
- What to expect during your recovery
- How to maximize your recovery
- Questions and answers
- Activity following your surgery

*Feel free to ask any questions you may have.
The more you know, the better prepared you will feel.*

PLEASE NOTE: Depending on your progress, you may be going directly home the day of or the day after surgery.

The importance of your coach

Friends and family are a major part of everyone's life and during this experience, their involvement is very important. We encourage you to choose a family member or close friend to act as your coach as you go through the hip surgery process. The coach works with you the entire time; from pre-op preparation, to inpatient recovery and into your discharge to home. Their help and support will make your journey easier.

Here are the things your coach can do to help you through your hip surgery experience.

BEFORE SURGERY

- Attend the pre-op education class with you.
- Prepare for your return home.
- Complete the pre-op checklist on the next page.

AT THE HOSPITAL

- Offer support and encouragement during exercise.
- Keep your morale high by sharing time and doing things that you like (board games, watching movies, etc.).
- Encourage you to participate in planned activities.
- Keep you focused on a healthy lifestyle and recovery.

AT HOME AFTER DISCHARGE

- Make sure you do the exercises. **No Exceptions!**
- See that you use assistive devices recommended to you as directed.
- Increase your activity level and to do things gradually as you gain your strength back.
- Oversee that you are following post-op instructions.
- Prepare healthy meals rich in vitamin C, calcium and iron.



Recovery Tip!

We encourage you to choose a family member or close friend to act as your coach as you go through the joint replacement process.

pre-operative HOME CHECKLIST

Preparing for your homecoming prior to your surgery will make your post-op days go smoother. Being prepared is the key to a relaxed recovery. Complete the list below.

- Make arrangements to have someone stay with you until you are comfortable being on your own. You will need help with bathing, dressing, meals and medications for a few days. Have enough food on hand or arrange to have someone go shopping for you.
- Do the laundry, change the linens etc., before leaving for the hospital.
- Bring a pair of comfortable shoes with non-skid soles for the hospital.
- Have easy access to a bed and bathroom on the downstairs level.
- Install a handrail if possible for any steps you may take routinely.
- Remove any obstacles that may cause you to trip: throw rugs, extension cords, low hanging bedspreads, pots, toys, pet toys etc.
- Make arrangements for your pets, mowing the lawn and bringing in the mail.
- Fill prescriptions for pain medication as well as routine prescriptions. Stool softeners, laxatives, Extra-Strength Tylenol® are examples of other medications you may need.
- Take care of financial matters such as bill payments, having cash on hand, etc.
- Arrange your plates, cups and utensils within easy reach to avoid using a step stool.
- Plan how you will transport your food to the table.
- Have a phone within easy reach with emergency numbers handy.
- Have a comfortable chair or couch with arms to help when getting up.
- Orthopedic surgeons require that you have your teeth cleaned prior to having a joint replacement.
- Have a full tank of gas before leaving for the hospital.



The hospital

WHAT YOU SHOULD BRING

We suggest that you bring some comfortable bed clothes, including a robe, pajamas or nightgown and slippers. You are welcome to bring your own toiletries. We will have some items for you if you need them. You may also want to bring your own books and magazines.

The most important item you should bring with you to the hospital is a list of your current medications. Be sure to include the current name, dosage and when you take them. Please include any over-the-counter medications, herbal supplements and vitamins. Be prepared to share this information when the Navigator calls.

VALUABLES

The hospital does not assume responsibility for valuables. This includes jewelry, dentures, hearing aids, watches, wallets, purses, eyeglasses, contact lenses and other personal items. Cell phones are particularly vulnerable to loss. Please make sure your coach has control of your cell phone at all time. They are small and easily left in sheets. If you should need items such as eyeglasses and hearing aids, keep them with you. When you go for surgery hand them over to a family member or your coach to hold or deposit them in the hospital safe. Money is best left at home.

PATIENT BILL OF RIGHTS

The law requires that every healthcare provider or facility provide you with a copy of the “Patient Bill of Rights and Responsibilities.” These rights and responsibilities are reviewed with you upon admission to the hospital, and signed by you and the admitting representative.

INFORMED CONSENT

As a patient, you have the right to informed participation in decisions involving your healthcare. This participation should be based on a clear, concise explanation of your condition and of all proposed technical procedures. This would include any risks, side effects, problems related to recuperation, probability of success or mortality. You should not undergo any procedure without your voluntary, competent understanding of the consent or the consent of your legally authorized representative.

pre-operative CHECKLIST



ROLE OF THE SPINE AND JOINT INSTITUTE NAVIGATOR

The Navigator

The Navigator will be responsible for managing your care from the pre-operative course through discharge including after discharge follow-up.

The Navigator will help you:

- Assess and plan for your return home, as well as your coach availability.
- Assess and plan for your specific care needs, such as anesthesia and medical clearance for surgery.
- Coordinate your discharge to outpatient services, home or skilled facility in the unlikely event continued care is needed.
- Assist you in getting answers to insurance questions.
- Act as your liaison throughout the course of treatment from pre-op through your discharge.
- Answer questions and coordinate your hospital care.

Shortly after your surgeon's office has scheduled your hip surgery, you will be contacted by the Navigator who will:

- Confirm your scheduled pre-op class and check that your pre-op medical testing is complete.
- Act as a liaison for coordination of your pre-op care between the doctor's offices, the hospital and testing facilities as necessary.
- Verify that you have made an appointment, if necessary, with your medical doctor and verify that you have obtained the pre-op tests your doctor has ordered.
- Answer questions and direct you to specific resources within the hospital including admissions or other departments as needed.

You may call the Navigator at anytime with questions or concerns about your upcoming surgery. **The Navigator's office is open from 9:30 am until 6:00 pm. (909) 793-4383.**

health QUESTIONNAIRE

Use this tool to gather your medical history in preparation for the pre-op assessment call.

1. Height and weight.
2. Any advanced directive (document stating your medical wishes), medical decision maker emergency contact name and phone number.
3. Privacy password: (needed by family/friends to obtain information about you).
4. Anesthesia/Drugs/Latex/Food/Environmental allergies and reactions.
5. Any over-the-counter medication including herbs, supplements, diet pills, vitamins (ex. Aspirin/Motrin, fish oil, ginkgo biloba).
6. Prescription medications (dosage & frequency).
7. Previous surgeries and hospitalizations including year of occurrence.
8. Exercise regimen and can you climb a flight of stairs without chest pain?
9. CONDITIONS: (including duration, frequency, and history of)
 - A. Heart
 - Heart attacks/chest pains/shortness of breath/dizziness
 - Heart surgery/angioplasty/stents/pacemaker
 - Irregular heart rate/palpitations/heart murmur/mitral valve proplase
 - Congestive Heart Failure/Swollen Ankles
 - High blood pressure
 - Blood clots/pulmonary embolism
 - B. Respiratory
 - COPD/asthma/emphysema/bronchitis
 - Pneumonia/tuberculosis
 - Lung cancer/cystic fibrosis
 - Sleep apnea/CPAP, BIBPAP use/snoring while asleep
 - C. Urinary
 - Renal failure/dialysis

health QUESTIONNAIRE

- Kidney stones/bladder tumors
- Prostate trouble
- Urinary frequency/urgency/incontinence/urinary tract infections
- D. Gastrointestinal**
 - Hepatitis/cirrhosis/pancreatitis/diverticulitis
 - Abdominal pain/constipation/diarrhea/bowel obstruction
 - Gastric reflux/hernia/ulcers
- E. Diabetes/thyroid**
- F. Central nervous system**
 - Seizures/strokes/numbness/tingling/headaches
 - Dementia/alzheimer's/confusion
- G. Blindness/glaucoma**
- H. Musculoskeletal**
 - Gout/arthritis/contractures
 - Use of assistive devices
- I. Skin conditions**
 - CDIFF/MRSA/VRE
 - Open wounds/lacerations
 - Eczema/psoriasis/shingles/rash
- J. Behavioral disorders**
 - Bipolar/schizophrenia/depression/suicidal thoughts/anxiety
- K. Date of last influenza and pneumonia vaccines**
- L. Tobacco/caffeine/alcohol/drug use & frequency**
- M. Pain control**
 - Frequency and duration of chronic pain
 - Effectiveness of pain medications
- 10. Pain control while in hospital**
 - A. Use pain control scale 0 – 10 (0=no pain, 10=worst pain)**

health QUESTIONNAIRE

- 10. Pain control while in hospital (continued)**
 - B. Pain control goal after surgery**
 - Your tolerable functioning pain level (Some pain is expected after surgery. Realistically, your pain will not be zero. We will make you as comfortable as possible.)
- 11. Discharge planning**
 - A. Do you have a coach?**
 - Assists during hospital stay/with home care and transportation
 - B. Any medical equipment needed?**
- 12. You will be asked to participate in a survey about your physicians at the Spine and Joint Institute at Redlands Community Hospital.**

The study which you are about to participate in is designed to examine your views about your health before your surgery; there is no right or wrong answers to this survey. You will also be called in 6 months and one year post-op to participate in the same survey. This helps us to see how you are healing after your surgery and how your quality of life has improved.

Please use this area to write down any questions, notes, concerns, etc. We look forward to speaking with you. Please have the answers to these questions available for your pre-op telephone assessment. We look forward to speaking with you.

Together, at the Spine and Joint Institute, we are committed to providing exemplary care in an encouraging, healing and rejuvenating atmosphere. Enthusiastically, we motivate and nurture your recovery with integrity and coordination with our patients, family and team members.

Thank you from the SJI team.

four weeks BEFORE SURGERY

START SUPPLEMENTS

Prior to your surgery, your surgeon may instruct you to start taking supplements to build up your body's healing mechanics. Typically, these are multivitamins with iron or with the iron taken separately.

READ "FREQUENTLY ASKED QUESTIONS" (FAQS)

Learn more about your surgery and what you can do to speed your recovery by reading the "Frequently Asked Questions" section.

REVIEW ADVANCED DIRECTIVES

In the event that you are unable to speak for yourself, having an advanced directive will ensure that your wishes concerning your healthcare decisions are honored.

Advanced directives require some thought, as well as paperwork. Prior to your surgery, the hospital will explain your options and assist you with completing the appropriate paperwork. Advanced directives are optional and require your signature in order to be enforced.

Please review the Advanced Directives section in this book on page 49.

ten days BEFORE SURGERY

STOP MEDICATIONS THAT INCREASE BLEEDING

Several medications, over-the-counter pain medications and even some vitamins may increase bleeding. Stop all anti-inflammatory and other medications that can cause bleeding 10 days prior to surgery. This includes aspirin, Motrin, naproxen, glucosamine, chondroitin, MSM, vitamin E, etc. If you need pain relief during his period, you can use Extra-Strength Tylenol®.



Discuss all medications with your doctor. Do not stop other prescribed medications without your doctor's direction.

Alert your doctor if you are currently taking a blood thinner like Coumadin, Lovenox, Plavix, or aspirin. Your doctor will give you special dosing instructions for stopping these medications.

three to five days BEFORE SURGERY

PREPARING FOR YOUR RETURN HOME

The first week or two after surgery, you may want to take it easy. It is best to have our house clean, the laundry done, the linens changed, monthly bills paid, etc., before you leave for the hospital. That way, when you come home, you can relax and focus on getting back into shape.

- **Stock up on nonperishable food, including pet food** if you have an animal.
- After surgery, you may be provided with a cooling unit to take home. Our staff will show you how to use the cooling unit before you are discharged. You can also use an ice pack. You can use a re-sealable bag filled with ice or any commercial ice bag. A bag of frozen peas works well because the bag readily conforms to any part of the body. The bag of peas can be refrozen and used repeatedly and then discarded when you no longer need it.
- **Prepare several meals in advance and freeze them.** That will make meal preparation after your return much easier. Likewise, arrange to have someone cut your grass, walk the dog, bring in your mail, etc.
- **Check your house for possible fall risks and remove any hazards.** Household items such as throw rugs and electrical cords in walkways that can cause you to trip.
- If you have not done so already, **place a rubber mat or non-skid adhesive strips on the bottom of your tub or shower.** Rent or purchase a bath seat if needed.
- **Remove any obstacles that may cause you to trip:** throw rugs, extension cords, low hanging bedspreads, pots, toys, pet toys etc.
- **Make sure that items you will be using are within easy reach.** Using a step stool or reaching a bottom cabinet may be a challenge when you first come home.

PRESCRIPTIONS

If you are currently taking medications, be sure you have an adequate supply to take you through the first few weeks. Fill any prescriptions the surgeon gives you for pain medication or blood thinners before you go to the hospital.

Have some Extra-Strength Tylenol® on hand, as well as stool softeners or laxatives.

the night BEFORE SURGERY

DO NOT EAT OR DRINK AFTER MIDNIGHT

Have a nice relaxing dinner the night before your surgery. Take any recommended medications. However, it is important that you **do not have anything to eat or drink after midnight**. This includes water, chewing gum and medications. *Eating or drinking after the recommended time could postpone your surgery. Please review the ERAS (Early Recovery After Surgery) handout and review instructions for pre-operative nutrition clear drink program with the Nurse Navigator.*

Please make sure to tell the Navigator if you are diabetic and you will be instructed accordingly.

If you normally take morning medications, wait until you get to the hospital. The pre-op nurse will give you instructions.

PACK THE FOLLOWING FOR THE HOSPITAL

- Steps to Success Guidebook
- Your email address and password for the Patient Wellbeing Survey
- Your insurance card
- Any co-payment required by your insurance company
- A driver's license or other photo ID
- Contact phone numbers
- Personal hygiene items
- Comb, brush, makeup, tooth brush
- Battery operated razor (no electric razors)
- Underwear and socks
- Flat shoes or tennis shoes
- Robe and sleepwear

BEST TO LEAVE A HOME

For your protection, please **do not bring** the following:

- √ Electrical items
- √ Valuables
- √ Jewelry
- √ Large amounts of money



Follow ERAS Guidelines

Do NOT eat or drink after midnight. Eating or drinking after midnight could postpone your surgery. If you normally take medications in the morning WAIT until you get to the hospital. The nurse will give you instructions.

Day of surgery

PRE-OP PREPARATION

Once you have been checked-in, you will be directed to the waiting area. You will speak to your anesthesiologist and operating room nurse as well as your surgeon. They can answer any last minute questions you may have.

As part of the surgical preparations, you will be given a surgical gown to wear and asked to remove any make up. It is not necessary to remove nail polish. After confirming which hip is to be operated on, the nurse may remove any hair in the surgical area, thoroughly scrub your hip with an antibacterial solution and start an IV. The use of an IV is necessary to administer fluids and medications during and after your surgery.

A Foley catheter may also be inserted to collect your urine during and for a short time after surgery.

MOVE TO RECOVERY

Immediately following surgery, you will be taken to the recovery room or Post-Anesthesia Care Unit (PACU). You will remain in the PACU for approximately ninety minutes after surgery while the anesthesia wears off.

- Your blood pressure and other vital signs will be closely monitored
- Pain control measures will be started
- Your doctor may have an x-ray taken of your new hip

SURGICAL WAITING AREA

While you are in surgery, your family and loved ones will be asked to wait in a waiting area. Once you are out of surgery, the surgeon will contact them and let them know your status. It is our goal to move you to your room within 90 minutes of your surgery finishing. Your family is able to wait for you in your room.



Recovery Tip!

Most people are brought directly from recovery to their rooms. Prepare to get started with your therapy activity within 60-90 minutes of being settled into your room. You may start activity in the PACU depending on the time of day. You may have visitors the day of surgery; however limit your visitors to one or two close friends or family members.

IN YOUR ROOM

Most people are brought directly from recovery to their room.

- Prepare to start your rehab activity as soon as you are fully awake from your anesthesia.
- The physical therapist will review how to do your exercises. These exercises are important; do them routinely.
- The nurse will be periodically checking your circulation and apply ice to your incision and help manage your pain.

one day AFTER SURGERY

Now it is time to get to work on your recovery. The staff will help you get set up for breakfast. You can also expect a visit from your surgeon, his assistant and/or medical doctor. The staff will help you get ready for your day and possibly ready for discharge.

Physical and Occupational Therapy will be done in your room today. They will help you to sit up for breakfast and you will begin work with the therapist. The therapist will begin focusing on activities of daily living to help you get ready for a smooth transition from the hospital to your home.

Medications for pain will be available at your request depending on your doctor's orders. It will be necessary to take pain medication. We want to plan the medications around your therapy sessions to avoid dizziness while trying to exercise. Switching to tablets rather than IV medications may be an option.

Dressings on your hip will remain in place. The doctor's request that the dressings are not removed from your surgical site unless they request it be changed. If you have drainage, we will reinforce your dressing.

CHANGE OF MEDICATION

Depending on your progress, IV pain medications will be replaced with oral medications. Oral medications are more effective than IV pain medications as they provide longer lasting pain relief.

PUT YOUR COACH TO WORK

Your coach is encouraged to be with you as much as possible during your hospital stay. Aside from keeping you company, this is an excellent opportunity for them to learn how to take care of you once you arrive home.

PLANNING YOUR DISCHARGE

Discharge time is 10 am.

Believe it or not, it's time to start planning on your return home.

Discharge planning



Discharge planning will start today. *Your Navigator will talk to you about your plans and needs for discharge.*

Excited about going home? Your nurse can help you with that!

DISCHARGE PLANS

Most hip surgery patients will be recovered enough to go home today. If you are not ready for discharge today, we will plan for tomorrow. Your Navigator will help you make your discharge arrangements. You will have therapy until time of discharge unless you have met all of your goals and your coach has taken over.

DISCHARGE DAY

Today is your discharge day. You will have therapy again today and it will focus on making sure you are able to get in and out of your car.

Your Navigator will help you make your discharge arrangements. Write down any questions that you may have so you can ask your doctor before you go home. Your doctor will write prescriptions for pain medications, if needed, to get filled on your way home or your doctor may have sent an electronic prescription to the pharmacy for you to pick up. The nurse will instruct you and your coach on the medications and side effects before you are discharged.

Your nurse will also give you written instructions for your medications, physical therapy, home equipment, follow-up appointments, contact numbers, restrictions and incision care. Please follow these instructions closely. If you have any questions once at home, do not hesitate to call with any concerns.

Most patients may go home by private car.



The Navigator will talk to you to assure your needs are met once you arrive home.

home CARE

You made great progress while you were in the hospital, but you are not fully recuperated yet. There is still work to be done.

GENERAL INFORMATION

- It is quite normal to feel tired and drained for a while after surgery. Do not let this get you down. It may take several weeks to get over this.
- When you can return to work is of course dependent on the type of work you do. Your doctor will discuss this with you.
- **No smoking, please. Smoking can increase the risk for surgical failure by 30%.**
- Make sure you elevate your legs twice a day for 30 minutes and wear your compression stockings as instructed to minimize swelling.

BE COMFORTABLE

It is human nature to avoid things that cause discomfort. If you are in pain, you are less likely to move or do your exercises. Inactivity can cause stiffness and will slow your recovery, undoing all your excellent work during your hospital stay.

- Take your medications 20-30 minutes before you begin your exercise at home or at physical therapy. This will make moving easier.
- Control discomfort by applying ice or using your ice equipment from the hospital. A bag of frozen peas works well because it will conform to any shape.
- Do not ice the surgical area for more than 15 minutes per hour. Non-stop ice can cause tissue damage and slow the healing process.
- Change your position every 45 minutes.

KEEP NAPPING TO A MINIMUM

During your recuperation, avoid napping during the day. This will allow you to sleep better at night.

eating right for FAST RECOVERY

How you treat your body the week before and after your surgery has a direct impact on how you heal. Now is the time to make the necessary preparations to give your body the building blocks it needs for a fast recovery. Don't wait for your surgery to be over – start the healing now.

- Exercise to increase muscle tone (See pre-op exercise section).
- Stop smoking.
- Avoid alcohol usage – especially 48 hours before surgery.
- Begin your healthy eating plan at least 10 days before surgery and continue for 10 days after surgery.
- Eat a well-balanced diet rich in iron, vitamin C and calcium.

On the following page are tables of iron, calcium and vitamin C rich foods you may want to add to your diet. You will also find a sample menu you can use as a guide.

HEALTHY EATING PLAN

Foods are divided into five basic food groups: vegetables, fruits, grains, milk and lean protein (meat, chicken and fish). **Unless you are on a specific diet prescribed by your doctor**, follow a daily well-balanced diet that includes:

- 2-3 servings of kale and orange vegetables such as carrots, squash and sweet potato
- 2 cups of fruit, but limit juices
- 6 ounces of grains: whole grain cereal, bread, crackers, brown rice
- 24 ounces of dairy: low-fat or skim, yogurt, low-fat cheeses
- 5-6 ounces lean protein and beans: lean meats, poultry, fish, pinto beans, lentils or kidney beans




IMPORTANT: Diabetics and people on other restricted diets should consult their doctor prior to starting this diet. People taking any type of prescribed medications, especially Coumadin and Plavix, should also check with their doctor before starting this diet. A sudden intake of increased green, leafy vegetables could react with your medication.

eating right for FAST RECOVERY

Try to incorporate as many high calcium, iron and vitamin C foods as you can into your diet.

A balanced diet and a good multivitamin will enhance the healing process. Ask your doctor which multivitamin is right for you and your medicine.



Iron is needed to build healthy red blood cells.

Vitamin C improves the body's absorption of iron.

A diet rich in calcium will help new bone regrow around your implant, making it more stable.



THE FOLLOWING IS A SAMPLE TO GET YOU STARTED:

BREAKFAST

- 1 egg cheese omelet with 1 slice low fat cheese, spinach and other vegetables
- 8 ounces of low-fat or fat-free milk/ may substitute milk with calcium added lactose free products
- 1 whole orange
- 1 piece of whole grain bread or toast

MORNING SNACK

- 1 cup of strawberries with low-fat plain yogurt

LUNCH

- 1 1/2 cups fresh spinach salad
- 2 ounces poultry (dark) baked, broiled or grilled

- 1/2 cup brown rice or 1 slice of whole grain bread
- 6 ounces low-fat or fat-free milk (may substitute milk with calcium added supplement)

AFTERNOON SNACK

- Green peppers with 2-3 ounces yogurt dip

DINNER

- 3 ounces lean meat, poultry or fish (baked, broiled or grilled)
- 1 medium baked potato
- 1/2 cup broccoli
- 1/2 cup of tomato

EVENING SNACK

- 1 cup low-fat or fat-free pudding or custard

painCONTROL

Pain control during and after surgery is one of the most common concerns of our replacement patients. With today's pain management techniques, you can be kept relatively comfortable.

ANESTHESIA DURING SURGERY

Anesthesia is the loss of sensitivity to pain brought about by various drugs known as anesthetics. There are several types to choose from. Your anesthesiologist will explain the one best suited for you.

General anesthesia is the most common form of anesthesia. The patient is put into a deep sleep and will not feel any sensation.

Spinal, regional or epidural anesthesia targets a specific area, like a joint, and totally numbs it. Although you are awake, you will not feel any pain. Typically with epidural anesthesia, another medication is administered to make you very relaxed and enter a light sleep state.

POST-OP PAIN CONTROL

There are several different kinds of pain control methods available that will keep you comfortable after surgery. Your doctor will choose the method right for you based upon your medical history, the amount of pain you are having and your phase of recovery.

- **Oral medications** are often used to control pain. These may also be administered prior to surgery to get a start on pain control and may be continued throughout your hospital stay. Most likely, you will be given a prescription for oral medication for use at home following discharge from the hospital.
- **Pain medications** are given in a multimodal approach and include narcotics and anti-inflammatories.

Regardless of the type of pain management being used, it is very important for you to communicate with your health care team if the pain medication is not sufficient, if you are nauseated or even if you are not as alert as you feel you should be. Adjustments can always be made to make you more comfortable.


PREVENT THE PAIN CYCLE

Pain has a cycle. It begins and increases until the pain medication interrupts it. The way to good pain control is to stop pain before it becomes intolerable.

If you begin to feel the pain increasing, please call your nurse. This is one time you do not want to "tough it out." Once the pain cycle takes hold, it will be very hard to control it.

PAIN SCALE

You will be asked to use a pain scale to help describe your pain level. If "0" means no pain and "10" is the worst pain possible, how would you rate the pain level? To relieve your pain most effectively, your healthcare team needs to know how well pain relief measures are working for you. Medications can be adjusted to meet your needs.



OTHER METHODS TO DECREASE PAIN

It is very important to relax after your surgical procedure. When you are relaxed, pain medications work better. You can also position yourself for comfort and ease of breathing. Applying ice to the area for 15-minute intervals may help.

Deep breathing can help relax tense muscles. Soft music can also help you relax.

Continue to use these pain control methods once you return home.



Recovery Tip!

Deep breathing can help relax tense muscles. Soft music can also help you relax. Continue to use these pain methods once your return home.

exercise your JOINT

Right now, your joint is probably somewhat stiff. Don't be concerned. You can remedy that with the right exercises. The following daily exercise routine will help restore flexibility and normal function to your new joint as well as guard against blood clots forming. Your doctor will have specific exercises. Please follow your therapists instructions.

Each day, as you do your exercises, you will notice that you can bend a bit easier than the day before. The key is to do the exercises faithfully as directed by your healthcare professional. A few minutes a day is a small price to pay for a pain free joint.

STANDING HIP ABDUCTION (Horizontal abduction/adduction)

1. Stand straight holding solid surface (counter or chair)
2. Toes pointed forward
3. Slowly kick sideways 10-12 inches
4. Do not lean sideways
5. Return to start position and repeat 10 times



STANDING HIP EXTENSION

1. Stand straight holding solid surface (counter or chair)
2. Toes pointed forward
3. Slowly kick backward 10-12 inches while squeezing glutes
4. Do not lean forward
5. Return to start position and repeat 10 times



exercise your JOINT


MINI SQUATS

1. Stand straight holding solid surface (counter or chair)
2. Bend hips and knees slightly as if to sit in a chair
3. Squeeze glutes to return to standing
4. Repeat 10 times



STANDING HIP FLEXION

1. Stand straight holding solid surface (counter or chair)
2. Slowly lift leg off floor flexing hip and knee
3. Return to start position repeat 10 times



exercise your JOINT


SINGLE LIMB ANKLE PUMPS

1. Lie on back with affected limb straight
2. Pull ankle back towards you as far as you can
3. Point toes as far as you can
4. Repeat 10 times



DOUBLE LIMB ANKLE PUMPS


1. Lie on back with both legs straight
2. Pull right ankle back towards you as far as you can and simultaneously point left toes as far as you can.
3. Alternate and repeat 10 times



exercise your JOINT


SUPINE HIP ABDUCTION

1. Lie flat on back with feet straight
2. Slowly slide/kick affected limb out to the side along bed
3. Slowly return to start position
4. Repeat 10 times



STRAIGHT LEG RAISE EXERCISE

1. Begin lying on back with affected leg straight
2. Slowly lift leg off surface keeping knee straight
3. Slowly return to start position
4. Repeat 10 times



GLUTE SETS

1. Begin at rest with legs straight
2. Squeeze buttocks and hold 5 seconds
3. Relax muscles and repeat 10 times



WEAN YOURSELF FROM PAIN MEDICATION

By the time you get home from the hospital, you will notice your need for pain medication decreasing. When you think you are ready, try substituting Extra-Strength Tylenol® in place of one dose of narcotic pain medication. Gradually increase the number of substitutions until you are no longer taking narcotics. Make sure you do not take more than 4 grams or 4,000mg of Extra-Strength Tylenol® in a 24 hour period.

If you are taking a blood thinner, check with your doctor prior to taking any other type of pain relievers. Many common, over-the-counter pain relievers may interact with your blood thinner and cause problems.



Patient portal

Access to your medical records anytime, anywhere! “My Health Info” provides you with the access to review and print lab results, discharge instructions, operative reports and more. Access to your health information is important to you. By using “My Health Info” you have your hospital history at your fingertips. You can access health information, update personal information, view upcoming appointments and access it all from your computer, smartphone, or tablet. If you would like to sign up, please visit our website at www.redlandshospital.org or call **Medical Records @ 909.335.5602**.

PREVENT CONSTIPATION

Changes in your daily routine, as well as taking narcotic pain medicines, can result in constipation. Take measures to prevent constipation before it becomes a problem.

- Eat fiber rich foods like grains, fresh fruits and vegetables to help keep your system moving.
- Drink plenty of water. This adds fluid to the colon and bulk to the stools, making bowel movements softer and easier to pass.
- Avoid liquids that contain caffeine, such as coffee and cola drinks. Caffeine flushes your colon of fluids and causes dehydration. Stools become dryer and harder to pass.
- Avoid alcohol. It also causes dehydration and contributes to constipation.
- Incorporate a daily walk or two into your exercise routine to keep your system well-balanced.
- Wean yourself from narcotic medications as soon as possible.

In the event you do become constipated, use stool softeners or laxatives such as milk of magnesia, as necessary.

BLOOD CLOTS AND ANTICOAGULANTS

Blood clots after surgery can cause serious problems. However, there are steps that can be taken to prevent them.

- Adhere to daily exercise
- Incorporate a daily walk or two

What to do if you suspect blood clots in the legs? By exercising, wearing the compression stockings as prescribed and taking your anticoagulation medication faithfully per your doctor’s instructions, your chance of developing a blood clot is minimal. However, it can happen. Prompt treatment usually prevents further complications.

- Call your surgeon IMMEDIATELY if there is swelling in your thigh, calf or ankle that does not decrease if you lie down with your feet elevated above heart level.
- Contact your surgeon if there is pain and tenderness in the calf of EITHER leg. DO NOT take a “wait and see if it gets better” attitude. If a clot occurs, it may be necessary to be admitted to the hospital to receive intravenous blood thinners for a short period of time.



If a clot occurs, it may be necessary to be admitted to the hospital to receive intravenous blood thinners for a short period of time.

Pulmonary embolus A pulmonary embolus is a blood clot that breaks away from the vein and travels through the blood into the lungs. This can be life threatening!

• **CALL 911 IMMEDIATELY** if you experience sudden chest pain, difficult or rapid breathing, shortness of breath, sweating or confusion. **DO NOT take the time to call your surgeon.**

The best way to avoid a pulmonary embolus is to recognize and treat any potential blood clots. **If you suspect a blood clot, call your surgeon IMMEDIATELY.**

PHYSICAL THERAPY

Your full participation in physical therapy is an essential element in the success of your hip replacement. The physical therapist will teach you the exercises your doctor has approved. Your Navigator will be in to visit you to help you arrange for any assistance that you may need at home.

PAIN

Pain is common and is to be expected after this type of surgery. Medication will be prescribed for you. Take as directed. Oral pain medication may cause nausea, constipation and a light-headed sensation. You should not drink alcohol or drive while on this medication. You may switch to over-the-counter pain medication such as Extra-Strength Tylenol® when your pain subsides.

INCISION

You should keep your dressing dry and in place until after your follow-up appointment with your surgeon, unless your surgeon has directed you otherwise. If you note any redness, swelling or drainage from your wound, please call your surgeon. You may be able to shower until you return for your follow-up appointment. The incision has not healed yet and getting the incision wet puts you at risk for infection. Do not go into a tub, swimming pool or jacuzzi to soak the wound for at least a month.

TEMPERATURE/FEVER

Your temperature may be slightly elevated for several days after your surgery. However, if fever persists above 101°F and is accompanied by chills, sweats, increased pain or drainage, you should call your doctor. These may be signs of infection. Make sure you do your deep breathing exercises and you can even use your incentive spirometer to help achieve full lung expansion. Deep breathing with the incentive spirometer also helps maintain core temperatures and helps with fevers too.

Swelling is common post-op. Normal swelling is reduced in the morning and gradually accumulates throughout the day. If the swelling is severe in the morning when waking you should contact your surgeon.



Precautions in positioning and movement at home

Avoid hip flexion greater than 90 degrees (a right angle).

Use your crutches or walker until your doctor tells you it is OK to stop using them.

Keep your feet and knees pointed straight ahead, not turned in or out.

Take small steps when you are turning. Try not to pivot.

Wear shoes with nonskid soles. Avoid wearing slippers as they can make you fall. Go slowly when you are walking on wet surfaces or uneven ground.

DO NOT dress standing up. Sit on a chair or the edge of your bed, if it is stable.

Try not to sit in the same position for more than 30 to 40 minutes at a time.

DO NOT sleep on the side of your new hip or on your stomach. If you are sleeping on your other side, place a pillow between your thighs.

ANTIBIOTICS

Prior to any dental, urological, gastrointestinal or surgical procedure, you must notify your doctor that you have had a joint replacement. You may need to take antibiotics to protect your joint replacement from infection.

PREVENTING COMPLICATIONS

As with all surgical procedures, there is a risk of complications such as deep vein thrombosis (DVT) or blood clot, atelectasis of lungs and infection.

DVT: While in the hospital, if the patient has a risk of DVT or a high risk medical history, the surgeon may also order anticoagulation medications. To prevent DVT, you should be out of bed on the first post-op day. You should initially ambulate with assistance.

Atelectasis or partial collapse of the lungs: You will be encouraged to deep breathe and cough and to use an incentive spirometer to decrease the risk of post-op respiratory issues. Early mobilization and ambulation also increases deep breathing and lung ventilation.

Infection: After discharge, you will be asked to notify your doctor if you develop new redness, swelling or drainage from your wound. It is important to keep your incision clean, dry and follow discharge instructions regarding the care of your incision. Also, notify your doctor if you have a fever above 101° and increase in pain.

MEDICATIONS AFTER DISCHARGE

Patients who go home after discharge will need to get their prescriptions filled soon after discharge, or better yet, prior to surgery.

PAIN PILLS

Pain pills may be needed for the first few days after surgery. You will notice, however, your need for narcotic pain medication will be decreasing. You can substitute Extra-Strength Tylenol® when you think you are ready to stop prescription pain medication. If you are taking a blood thinner, check with your doctor before taking any other type of over-the-counter pain medication since some may increase the effects of blood thinners.

Remember that many prescription pain medications can cause constipation. Take measures to prevent constipation such as:

- Drink plenty of water which adds fluid to the colon, making bowel movements softer and easier to pass.
- Eat fiber rich foods like grains, fresh fruits and vegetables to help keep your system moving.
- Avoid liquids that contain caffeine such as coffee, and cola drinks which cause dehydration.

- Avoid alcohol which also causes dehydration.
- Try to incorporate a daily walk into your exercise routine to keep your system well balanced.

By exercising and taking blood thinners (if your doctor deems necessary), your chance of developing a blood clot is minimal. However, if you notice these symptoms, please call your doctor.

Prompt treatment can prevent further complications:

- If you notice swelling of your thigh, calf or ankle, lie down with your feet elevated above the level of your heart. If the swelling does not go down, call your doctor immediately and do not put any weight on the swollen leg. Avoid rubbing your leg even if it's painful.
- **Contact your doctor if there is pain in the calf of EITHER leg.** DO NOT take a “wait and see” attitude if you have pain in your calf.
- If you suspect a blood clot, call your surgeon immediately.

If a blood clot occurs, it may be necessary to be admitted to the hospital to receive intravenous (IV) blood thinners for a short period of time.



If admitted to the hospital, DO NOT TAKE THE TIME TO CALL YOUR DOCTOR. The hospital will notify your doctor once you are admitted. The best way to avoid a pulmonary embolus is to recognize and treat any potential blood clot.

ASPIRIN

Ongoing studies indicate that aspirin may be useful in preventing blood clots after surgery. Because aspirin has anticoagulant properties, it should not be used when taking other anticoagulants.

Your nurse will tell you how much aspirin to take if your doctor has prescribed it.

REPORT ANY ANTICOAGULANT RELATED SIDE EFFECTS IMMEDIATELY

If taken according to your surgeon's instructions, anticoagulants are safe and effective. However, in some cases, there may be warning signs that require prompt treatment.

home CARE

If you fall, have a traumatic injury or if you experience any of the following, call your surgeon immediately:

- Bleeding or oozing from surgical wound
- Bleeding at the site of injection
- Nosebleeds
- Blood in your urine
- Coughing or vomiting blood
- Excessive bleeding when brushing your teeth
- Spontaneous bruising (not caused by a blow or any apparent reason)
- Pain or swelling of any part of your leg, foot or hip
- Dizziness, numbness or tingling
- Rapid or unusual heartbeat
- Chest pain or shortness of breath
- Vomiting, nausea or fever
- Confusion

PREVENTING SURGICAL SITE INFECTIONS

Infections, although rare, do sometimes occur after surgery. It is important to note any changes in your incision.

SIGNS OF INFECTION

- Some redness, swelling and bruising around the incision are perfectly normal. Call your doctor/surgeon if the redness increases or the pain doesn't subside.
- Report any fever or night sweats to your doctor/surgeon.
- Call your doctor/surgeon if you notice any increase in drainage, if the clear discharge changes color or if an odor is present.
- Call your doctor/surgeon if you notice an increase in pain (not associated with normal exercise).

home CARE

PREVENTING INFECTION

The first step: Taking proper care of your incision and dressing.

The second step: Notify your healthcare provider prior to any procedure that may break the skin. This includes dental procedures.



Your doctor will give you specific instructions for the care of your dressing before you are discharged from the hospital.



GENERAL INSTRUCTIONS INCLUDE:

- Keep the incision clean and dry.
- Do not use any lotions, rubs, ointments, etc. on your wound unless directed to do so by your home health care nurse or doctor.
- Examine your wound daily when dressing changes are done and report any sign of infection to your doctor.
- Do not get the incision wet in the shower until instructed by your doctor. Do not immerse under water until after seen on follow-up appointment with your doctor
- You may have a water resistant dressing.
- Inspect the wound for any signs of infection such as increased swelling, redness, yellow or green discharge and odor. Report any abnormal findings to your surgeon.

home CARE

- Swab the incision area with Betadine antiseptic solution, if desired.
- Pick up the dressing pad by the corner. Do not touch the part of the pad that is laid over the incision. This could introduce bacteria into the wound and cause an infection.

Recovery Tip!

Take a proactive approach to prevent infection. Taking Proper care of your incision is the first step. The second Step is to notify your healthcare provider prior to any Procedure which may break the skin. This also applies to any dental procedures.



everyday LIVING

FOLLOW-UP VISITS

Keep your scheduled post-op visit. If you do not have one, please call your doctor's office to schedule it.

SELF-CARE

The best thing to do is to think before you act. Ask yourself if the activity you are about to perform can be done so safely. Taking a moment to assess the situation will also make you more aware of the alignment of your body and arm. Allow plenty of extra time for normal, daily activities. Do not hesitate to ask for assistance in performing any activity, even if it seems simple. Let your family, friends and co-workers be a part of your healing. Remember, they know of your procedure and want the best possible outcome for you.

BATHING

Make sure all surfaces are dry before you get into the tub or shower. Walk-in showers are recommended over baths because they are easier to get in and out of. Long-handled bath sponges and hand-held shower hoses are helpful. If showering, hang a shower caddy over the shower nozzle to keep your bath articles within easy reach. You can place a waterproof armchair in the shower for safety. If possible, have your coach supervise you during your first few times getting in and out of the shower/tub. Keep in mind that you might want a rubber mat on the floor of the shower or the tub to help protect against slipping.

TOILET

During the first few weeks after surgery, you may need extra time to complete toileting tasks.

RESTING AND SLEEPING

Use a firm mattress or couch. Soft pillows provide support for your neck and legs. Sleeping promotes healing, it is important that you get sufficient rest as you recuperate. As you rest, you may become stiff, so get out of bed slowly.

DRESSING

Allow yourself plenty of extra time to dress. Hurrying may result in improper arm motions. Loose fitting clothes, such as sweatpants or shorts with elastic waists, and slip on shoes are recommended. Your therapist might recommend using a sock aid. Carefully dress using a chair for support and to protect against loss of balance while you are not able to use your surgical arm.

SITTING

When sitting, carefully reach for the chair prior to lowering your body onto the chair. This will insure that the chair does not move as you sit down.

STANDING

Make sure that you keep your posture upright to maintain a natural standing posture.

Call 911 Immediately if you experience sudden chest pain, difficult or rapid breathing, shortness of breath, sweating or confusion. **DO NOT** take the time to call your surgeon.

PLEASE NOTE: If you fall or have a traumatic injury, call your surgeon immediately.

KITCHEN ACTIVITY

Whenever possible, keep frequently used items on the counter top. This includes pots, pans, bowls, storage containers and spices. You may not like the way the counter looks, but it will be back to normal soon. The use of a reacher is highly recommended to pickup items either above or below the counter. Do not hesitate to obtain assistance at the store and at home.

LAUNDRY

If using a top or front-loading machine, stand close to the washer and maintain your balance. Small loads are much easier on you than large loads. Wet clothes weigh much more, so remove them slowly, no more than one or two items at a time.

LIFTING

Avoid lifting as much as possible and ask for assistance. Do not lift objects that are awkward or weigh more than 5 pounds until your doctor says you may lift more. Move slowly and avoid sudden, jerky movements. Be sure you test the weight of the object prior to attempting to move it.

DRIVING/RIDING IN A CAR

Do not drive until you receive your surgeon's approval. You may want to add a firm pillow on the seat to make it higher. You may also incline the backrest and push the car seat backwards in order to have more space to get in and out of the vehicle. When getting in, sit first, and then slowly swing both your legs in at the same time.



PREVENTING WEIGHT GAIN

Weight gain is not uncommon after surgery. Between decreased activities, boredom associated with limited mobility, depression and pain, those extra pounds can creep up on you before you know it. Unfortunately, excess weight only fuels the vicious cycle of increased pain, decreased activity and depression.

These suggestions may be helpful in preventing excessive eating and weight gain after surgery:

- **Drink at least two quarts of water each day.** This will help you to feel full and you will be less likely to overeat.
- **Fill your refrigerator and cupboards with healthy foods.** The internet and library are endless resources for healthy eating habits and low-fat recipes.
- **Chew foods with a lot of “chewing power”.** It actually takes 20 minutes for your stomach to tell your brain it is full. If you are chewing a long time, you will be satisfied with smaller portions. Healthy foods with “chewing power” include:
 - * Popcorn (easy on the butter and salt)
 - * Pretzels
 - * Apples
 - * Celery
 - * Carrots
 - * Bagels
 - * Chewing gum also works.

For example, picture how many apples you can eat in 20 minutes. Now picture how much ice cream you can eat in 20 minutes.

Keep your mind and your hands busy. Watching TV for extended periods of time is not only bad for your hip recovery, but commercials often entice us to eat when we are not hungry. Focus on crossword puzzles, building models, needlepoint, knitting, card games, board games or anything else that will divert your attention away from TV and the refrigerator. This is a great opportunity to spend some quality time with family and friends.

Any kind of surgery carries a certain amount of risk. We will take every precaution to assure that your surgery is successful and without incident. However, in the event of an unexpected complication, we want to honor your wishes and individual considerations. It is very important to put your healthcare preferences in writing prior to your surgery.

Advanced directives are a means of directing your medical care in the event that you are unable to do so for yourself. Once the advanced directives are on file, the doctor, your family and hospital staff are committed to honoring your wishes. There are many forms that can be obtained either from the internet, i.e., “Five Wishes” or you can ask the Navigator to provide forms at orientation.

THREE TYPES OF ADVANCED DIRECTIVES


1. LIVING WILLS detail your wishes for healthcare if you have a terminal condition or irreversible coma and are unable to communicate. Normally, this refers to the level of life support measures you would like to have administered in order to prolong the dying process when death is eminent.

2. APPOINTMENT OF A HEALTHCARE AGENT is a process that authorizes a person of your choice to make medical decisions for you in the event you are unable to do so for yourself. It is more flexible than a living will because it can cover any healthcare decision, even if you are not terminally ill or permanently unconscious. This type of advanced directive is also referred to as a Medical Power of Attorney.

3. HEALTHCARE INSTRUCTIONS are your specific choices regarding use of life sustaining equipment, blood products, hydration, nutrition and the use of pain medications. You should also make your wishes known in terms of any organ donations.



Frequently ASKED QUESTIONS



Jot down any other questions you may have in the journal section of this workbook.

People who are facing joint replacement surgery typically ask the same questions. However, if you have questions that are not covered in this section, please ask your surgeon or anyone on the joint care team. We are here to help.

WHAT IS ARTHRITIS AND HOW DOES IT AFFECT MY HIP?

Arthritis is an inflammation of the bone that occurs when bone rubs against bone. In the hip, the ball or femoral head at the top of the thigh bone fits into a

cuplike area or socket located at the bottom of the pelvis or hip. A healthy hip normally has a layer of gelatin like material (cartilage), located between the ball and socket. The cartilage allows the ball to move smoothly inside the socket when you change positions.

Over time, the cartilage begins to wear away. With little or no cartilage to serve as a cushion, the nerve endings surrounding the ball and socket become irritated. Bone-on-bone contact is responsible for the pain, swelling and stiffness that are associated with arthritis.

Frequently ASKED QUESTIONS

WHAT IS TOTAL HIP REPLACEMENT?

The term total hip replacement is misleading. The hip is not totally removed, to be replaced by a new hip. Rather, the femoral head is removed and replaced with a metal ball. Likewise, the worn socket (acetabulum) at the base of the pelvis is reshaped and a cup-like device is inserted. Bone-on-bone contact is eliminated and so is the bone-on-bone pain.

WHAT IS PROSTHESIS?

The artificial ball and socket are referred to as the prosthesis or implant. There are several types of prostheses. Some are made totally of metal, with plastic liners in the socket while others may incorporate ceramic components. Your surgeon will determine which type of prosthesis is most suitable for you.

IS IT POSSIBLE TO HAVE BOTH HIPs DONE AT THE SAME TIME?

It is possible to have both of your hips done at the same time. This is called a bilateral total joint replacement.

Unilateral joint replacement (one hip at a time) is normally recommended. During the recuperation process it will be easier to walk and exercise if only one hip is undergoing the healing process. Usually, the second hip can be scheduled within three to four months of the original surgery. This will allow the first hip ample time to heal.

WHAT IS THE DIFFERENCE BETWEEN A CEMENTED OR UNCEMENTED PROSTHESIS?

See the Hip Replacement image on page 50.

When using the cemented technique, a prosthesis with a smooth finish is cemented or glued into place. In the uncemented technique, a prosthesis with a

porous coating is placed directly into the bone. It is held in place by the elasticity of the bone tissue and by basic friction between the bone and the prosthesis.

Over time, new bone will grow and attach to the porous coating anchoring the prosthesis firmly into place.

WHICH IS BETTER - CEMENTED OR UNCEMENTED PROSTHESIS?

The answer to this question is different for different people. Because each person's condition is unique, you and your doctor must weigh the advantages and disadvantages.

Cemented replacements are more frequently used for less active people and people with weaker bones or osteoporosis. Uncemented replacements are more frequently used for more active people. Studies show that cemented and uncemented prosthesis have comparable rates of success.

WHAT IS LAMINAR FLOW?

Laminar flow is a type of air circulation and filtration system used in some operating rooms. The system filters and circulates that air in the room removing airborne bacteria. Since bacteria can cause post-op infection, it was thought that the cleaner the air, the less chance of infection.

With improved surgical techniques, use of antibiotics plus multiple other infection control measures, post-op infection rates have decreased and, consequently, laminar flow rooms have become less popular.

WHAT IS THE SUCCESS RATE FOR TOTAL HIP REPLACEMENTS?

Approximately 90-95% of patients report good to excellent results in terms of pain relief. Most people are able to significantly increase their activity and mobility and return to the low impact activities they participated in prior to the onset of arthritis pain.

Frequently ASKED QUESTIONS

AM I TOO OLD FOR THIS SURGERY?

Your overall health is more of a determining factor than your age. Prior to the surgery, you will be asked to see your family doctor to assess any health risks. All measures will be taken to prepare you for a successful surgical outcome.

HOW LONG WILL MY NEW HIP LAST?

There are no guarantees on how long your new hip will last. Various factors such as weight, activity type, activity level, etc. can affect the usable life of your new hip prosthesis. Current studies indicate that the average hip prosthesis lasts for 15 to 20 years. With new materials and procedures, this expectation may change.

WILL I NEED TO HAVE MY HIP REPLACED AGAIN IN THE FUTURE?

Some people have a hip replacement that lasts their entire lives; other people need to have the procedure repeated. If the bone does not properly bond to the first replacement, the prosthesis becomes unstable and needs to be replaced. If the plastic spacer that cushions the joint becomes extremely worn, this may also require replacing.

ARE THERE MAJOR RISKS ASSOCIATED WITH THIS TYPE OF SURGERY?

All surgeries carry a certain amount of risk. However, because of our proactive approach in preventing possible complications, most of our joint patients are just fine and are ready to leave the hospital in a couple of days. We take special care to safeguard you from infection following surgery. To further minimize the risk of infection, we have streamlined the surgical procedure to take less time. The less time your wound is open, the less chance of infection.

Following surgery, blood clots can become a problem. You may be given medication to reduce the risk of blood clots forming. Your surgeon may prescribe an anticoagulant such as aspirin, Fragmin, Coumadin, Lovenox or Rivaroxaban. Getting you up and walking soon after surgery is another way to reduce the risk of blood clot.

HOW LONG DOES HIP REPLACEMENT SURGERY TAKE?

The surgery itself takes about one to two hours. After the surgery you will be monitored closely in a special unit called the Post Anesthesia Care Unit (PACU) until the anesthesia wears off. Once you are awake and stable, you will be transferred to your room.

WHO WILL BE DOING THE SURGERY?

Your orthopedic surgeon will be performing your surgery. Typically, an assistant will help during the procedure. You or your insurance may be billed separately for this assistant's services.

WILL I BE AWAKE DURING THE SURGERY?

Normally, during the surgery, an anesthesiologist will administer an anesthetic: a general anesthetic will put you into a deep sleep, while a regional anesthetic will numb specific areas only. In a normal situation, regional anesthetics are given with another medication that will make you very relaxed and put you into a light, dreamlike state. You will not remember the surgery once you awake.

You and your anesthesiologist will discuss which method is best for you prior to your surgery. Feel free to discuss any concerns you may have.



For any concerns about anesthesia read the Pain Control section of this book.

Frequently ASKED QUESTIONS

WHAT WILL MY SCAR LOOK LIKE?

Several different techniques are used for hip replacement surgery. The type of technique used will determine the number, location and length of the scar(s). The traditional approach is to make an incision running lengthwise over the side of the hip.

Minimally invasive surgeries are becoming more popular. Special instruments are used that allow the surgeon to replace the hip joint through one or two smaller incisions. Your surgeon will discuss which type of surgery is best for you.

There may be some numbness around the scar after it is healed. This is perfectly normal and should not cause any concern. The numbness usually disappears over time.

OTHER THAN THE SCAR, WILL MY JOINT BE ANY DIFFERENT?

Normally, the new joint feels completely natural. However, some people may notice that the leg with the new hip seems slightly longer than it was before. Years of slow degeneration or breakdown of the hip joint may have caused a shortening effect. This process can be so slow that your body will just learn to compensate for the difference. Once the joint components are replaced, the leg feels longer. In other instances, the surgeon will intentionally lengthen the leg at the joint during the surgery. This is done to prevent the joint from dislocating or popping out of the socket. As time passes, the increased leg length becomes less noticeable.

Sometimes a small lift placed inside the opposing shoe is needed to correct the length difference. Consult your surgeon if you encounter this problem.

A few patients report aching in the thigh when standing or weight bearing. Normally this aching will disappear after a few months and the hip is pain free, so patients are ready to resume an activity lifestyle once again.

WILL I BE IN A LOT OF PAIN?

Our joint center specializes in joint replacement surgeries. As such, we have considerable experience in caring for patients after surgery and know how to keep our patients comfortable.



For more information, read the Pain Control section of this book.



WILL I NEED A PRIVATE NURSE AFTER SURGERY?

There will be no need for a private nurse. Our joint center's care team are the experts.

WILL I NEED A BLOOD TRANSFUSION AFTER SURGERY?

Your blood will be typed up on admission. There is a possibility that you may need a blood transfusion after surgery. If you need a transfusion, your doctor will discuss it with you.

Frequently ASKED QUESTIONS

HOW LONG WILL I BE CONFINED TO BED AFTER SURGERY?

On the day of your surgery, the Physical Therapy team will evaluate you to see if it's safe to get you moving. Your care team will advise you when it is safe to get up. Early the next morning, you will be up and dressed to start the day's activity. Usually, most patients will be walking with a walker or crutches by the afternoon.

HOW LONG WILL I BE IN THE HOSPITAL?

Hip replacement patients usually stay with us one or two days; however, some may leave earlier. Before you leave, you must meet certain goals. You will learn more about these goals in your group exercise class.

WILL I NEED A WALKER, CRUTCHES OR CANE?

People progress at their own rate. Normally, patients use an assistive device such as a walker or cane for about four to six weeks. Your doctor will tell you when it is time for you to retire them.



CAN I GO DIRECTLY HOME, OR DO I HAVE TO GO TO A NURSING HOME/ REHAB CENTER?

Occasionally, some patients require a short stay of three to five days in a skilled nursing facility, but this is the exception rather than the rule. Your care team will be monitoring your progress on a daily basis and will determine if you are ready to go directly home.

WILL I NEED HELP AT HOME DURING THE FIRST WEEK?

Although you will be well on your way to recovery when you leave the hospital or skilled nursing facility, you will need to arrange for someone to assist you with your meals, medication, dressing, etc. at least for the first week or two. If you go directly home from the hospital, the joint care team can arrange for a home health care nurse to visit your home, as needed. Be sure to alert the joint care team if you live alone. This may or may not be covered by insurance.

To make the transition to home easier, plan ahead. Prior to coming to the hospital, take care of such things as getting prescriptions filled, changing the beds, doing the laundry, washing the floors, arranging for someone to cut the grass, walk the dog, stocking up on groceries, etc. Your job after surgery is to focus on your recovery, not household tasks.

WILL I NEED PHYSICAL THERAPY WHEN I GO HOME?

Most patients receive physical therapy after they are discharged from the hospital. Your surgeon's office will help coordinate your outpatient therapy.

The number of physical therapy sessions will be based on your individual progress. To a large extent, your progress will be determined by how much effort you put into your exercise routines. Instructions for your pre- and post-op exercises are included in this book.

Frequently ASKED QUESTIONS

WHY SHOULD I EXERCISE BEFORE SURGERY?

The better the condition your muscles are in prior to the surgery, the easier and faster your recuperation is expected to be. It is important to learn the exercises and be comfortable with them prior to the surgery so that you can continue them once you return home. Starting the exercises now will build muscle tone and pave the way to a quick recovery.

Begin doing the exercises immediately. Your new joint will be happy that you did.

AFTER LEAVING THE HOSPITAL, WHEN DO I NEED TO SEE MY SURGEON AGAIN?

You should plan a follow-up visit as scheduled with your surgeon. If you have questions about your follow-up appointment, please contact your surgeon's office immediately.

ARE THERE ANY ACTIVITIES THAT I SHOULD AVOID INITIALLY?

Keeping your new joint moving certainly will help your recovery process. However, you should return to your normal activities gradually. In some instances, you may have to work your way up to a particular activity. Taking a five mile walk on your first time out, for example, is not realistic. Rather, walk until you begin to get tired. Add distance to each subsequent walk until you have reached your goal.

You will be instructed by your joint care team to avoid specific positions of the joint that could lead to dislocation. Avoid high impact activities, such as jogging, singles tennis, basketball, down-hill skiing, football, etc. Consult your surgeon prior to participating in any high impact or injury-prone sports.

WHEN CAN I RESUME HAVING SEXUAL INTERCOURSE?

After surgery it will take time to regain your strength, as well as confidence in your new hip. Most people feel able, physically and mentally, to engage in sexual activity about four to six weeks after surgery.

Although individuals vary in their healing rate, at the four to six week point the incision, muscles and ligaments are usually sufficiently healed to consider resuming sexual activity. Talk to your surgeon if you have any questions.

WILL MY MEDICATIONS AFFECT MY ABILITY TO ENGAGE IN SEXUAL INTERCOURSE?

Some medications can affect your performance and/or enjoyment during intercourse. Many narcotic pain relievers and cortisone medications can decrease sexual performance. Other common medication-related side effects are a decrease interest in sex, abnormal erections, vaginal dryness, and delayed orgasms.

If you sense that your medication is causing these side effects, try having sex in the morning before taking your first dose or in the evening before your last dose.

DO NOT adjust or stop taking your prescribed medicine without consulting your surgeon. Normally, a simple adjustment or change in medication can eliminate unwanted side effects.



DO NOT ADJUST OR STOP TAKING YOUR PRESCRIBED MEDICINE WITHOUT CONSULTING WITH YOUR SURGEON. Normally, a simple adjustment or change in your medication can eliminate unwanted side effects.

Frequently ASKED QUESTIONS



ARE THERE ANY POSITIONS THAT SHOULD BE AVOIDED DURING SEXUAL INTERCOURSE?

Females should keep affected leg out to the side with one or two pillows under the thigh and knee for support. Move the affected leg as little as possible.

Later in the recovery process as the hip swelling decreases and range of motion improves, you may want to experiment with other positions. Stop if you experience discomfort. In case of severe pain, DO NOT MOVE. Call your surgeon immediately for instructions.

SIDE LAYING POSITION

- The female joint replacement patient should lie on the unaffected joint side, using at least two pillows between the legs to support the affected leg. Position the pillows to prevent the affected leg from crossing over the other leg.
- The male joint replacement patient should lie on the unaffected joint side and use his partner's legs to support his affected legs. The female partner should place at least two pillows between their legs so that the male's affected leg is positioned safely. Care

should be taken not to bend the hip over 90 degrees or let the leg cross over to the mattress.

TOP POSITION (SUGGESTED FOR MALES)

- When assuming the top position from a back lying position, DO NOT roll your affected leg inward towards the other leg. Keep it straight and in-line with the hip.
- DO NOT bend the affected hip more than 90 degrees.
- Keep the affected leg out to the side with the toes pointed slightly outward.

SITTING POSITION (SUGGESTED FOR MALES)

- Place one or two pillows on the bed or chair so that the hips are higher than the knees when sitting. The hip should be bent less than 90 degrees.
- DO NOT allow the knee of the affected leg to raise past the level of the navel.
- Be careful when exiting the chair or bed. DO NOT bend the affected hip more than 90 degrees.

WHAT IF SEXUAL INTERCOURSE DOESN'T GO WELL?

Remember, you are still in the healing process. Just like other activities that you are returning to, it may take some time to regain your former stamina. Realize that these changes to your sex life are temporary and are needed to protect your new hip joint. Just relax. You will be back to your old self in no time.

WHEN CAN I DRIVE?

How soon you can resume driving depends on several factors. It can be as little as four weeks or as long as six weeks or more. Your return to driving largely depends on how committed you are to your exercises and physical rehabilitation.

Frequently ASKED QUESTIONS

Another consideration is the mechanics it takes to drive your car. If you have had a left hip replaced, you may be able to drive a car with an automatic transmission in as little as four weeks depending upon your own personal progress. If your surgery was on your right hip or you are driving a car with manual transmission requiring both feet, then you may not be ready to drive for six or more weeks. It all depends how far your recovery has progressed. Regardless of your progress, you should not consider driving, especially if you are still taking prescription pain medication.



Your surgeon will let you know when it is advisable to drive again.

ARE ANY ACTIVITIES BETTER THAN OTHERS?

Exercise is very important to the entire body to maintain health. It is especially beneficial for your new joint.

Ask your doctor when it is safe for you to incorporate low impact activities such as golf, dancing, hiking, swimming, bowling, gardening, etc. back into your normal routine.

WHEN CAN I RETURN BACK TO WORK?

The physical demands required for your job, as well as your own progress, will determine when you can return to work. Typically, people plan on taking a one month leave of absence from work. Some people with sedentary jobs may be able to return sooner. Your surgeon will tell you when you can return to work.

WILL MY NEW HIP SET OFF SECURITY SENSORS WHEN TRAVELING?

The prosthesis is made of a metal alloy and may or may not be detected when going through some

security devices. If you wish to have a copy of your operative report, please ask a member of your care team to assist you with this.

WHAT ABOUT MY CURRENT MEDICATIONS?

Discuss your current medications with your physician to see if and when you should modify your medication schedule. Remember to include any over-the-counter drugs as well as vitamins and herbal supplements. It is important that your healthcare team know all of the medications, nutritional and herbal supplements that you are taking because some may cause problems such as excessive bleeding during surgery. This may or may not be relevant to you depending on a number of factors, including:

- The surgical procedure you are having.
- Your medical history.
- The medications/supplements you are taking.

It is very important to give accurate information to your healthcare team. Do not stop or change any of your medications unless instructed.

SHOULD I CONTINUE TO TAKE MY BLOOD PRESSURE MEDICATION?

If you take an oral diabetes medication such as glyburide, you may be told not to take it the day of surgery. Other oral medications such as Glucophage may be stopped for several days prior to surgery as this medication may cause serious side effects while under anesthesia.

Insulin will be continued, but usually at a different dose in order to prevent your blood sugar from dropping to low during fasting.

Frequently ASKED QUESTIONS

ANESTHESIA FAQS

WHO ARE THE ANESTHESIOLOGISTS?

The Operating Room, Post-Anesthesia Care Unit (PACU) and Intensive Care Units at the Spine and Joint Institute are staffed by board certified and board eligible physician anesthesiologists. Every anesthesiologist is an individual practitioner with privileges to practice at the Spine and Joint Institute.

DURING SURGERY, WHAT DOES THE ANESTHESIOLOGIST DO?

The anesthesiologist is responsible for your comfort and well-being before, during and immediately after your surgical procedure. In the operating room, the anesthesiologist will manage vital functions, including heart rate and rhythm, blood pressure, body temperature and breathing.

The anesthesiologist is also responsible for fluid and blood replacement, when necessary.

WILL I HAVE ANY SIDE EFFECTS?

The anesthesiologist will discuss the risks and benefits associated with the different anesthetic options, as well as any complications or side effects that can occur with each type of anesthetic. Nausea and vomiting may be related to anesthesia or the type of surgical procedure. Although less of a problem today because of improved anesthetic agents and techniques, these side effects continue to occur for some patients. Medications to treat nausea and vomiting will be given if needed. The amount of discomfort you experience will depend on several factors, especially the type of surgery.

Your doctors and nurses can relieve pain with medications. Your discomfort should be tolerable, but do not expect to be totally pain free.

WHAT WILL HAPPEN BEFORE MY SURGERY?

You will meet our anesthesiologist immediately before your surgery. The anesthesiologist will review all the information needed to assess your general health. This includes your medical history, laboratory test results, allergies and current medications. They will then determine the type of anesthesia best suited for you. He or she will also answer any additional questions you may have.

You will also meet your surgical nurses. Intravenous (IV) fluids will be started and pre-op medications may be given. Once in the operating room, monitoring devices, such as a blood pressure cuff, EKG and other devices will be attached to you for your safety. You are now ready for anesthesia.

WHAT CAN I EXPECT AFTER THE OPERATION?

After surgery, you will be taken to the PACU. You will be watched closely by specially trained nurses. During this time period your breathing and heart functions will be closely monitored and you may be given extra oxygen, if necessary.

An anesthesiologist is available to provide care as needed for your safe recovery.

my surgery JOURNAL

DAY OF SURGERY

DATE: _____

POST-OP: DAY ONE

DATE: _____

my surgery JOURNAL

POST-OP: DAY TWO

DATE: _____

POST-OP: DAY THREE

DATE: _____

my surgery JOURNAL

QUESTIONS OR CONCERNS
